

ginal design. It will contain a very full account of all that has been hitherto written on the subject; and an original figure of every species and remarkable variety will be given. Dr. Johnston intends at the same time to avail himself of this opportunity of adding many new figures of zoophytes contributed by his friends, and acquired by himself, since the volume on British Zoophytes was published.

*Corpus Florarum Provincialium Sueciæ.*—I. *Floram Scanicam scripsit* E. Fries, Upsaliæ, 1835. 8vo.

This is the first of a series of Floras of the different parts of Sweden, comprising that of Schonen. Our object in noticing it here is to give a sketch of the learned Professor's latest views on the arrangement of Fungi, which we are enabled to do not only by the help of the work itself, but from a letter from the Professor himself, dated October 4th, 1838. As the work (though, like the 'Novitiæ Floræ Suecicæ' lately noticed, it is of great utility) is not probably in many hands, we think it cannot fail to be acceptable.

1. HYMENOMYCETES. { *a.* Agaricinei. *b.* Polyporei. *c.* Hydnei. *d.* Auricularinei. *e.* Clavariæ. *f.* Tremellinei.
2. DISCOMYCETES. { *a.* Helvellei (Morchella, Helvella, Rhizina, Peziza, &c.) = Pez. ceracæ.  
*Discus excipulo saturatior.* { *b.* Bulgariæ (Leotia, Bulgaria, &c.) = Pez. gelatinosæ.  
*Discus ut in sequentibus excipulo dilutior.* { *c.* Dermei (Vibrissea, Ditiola, Dermea, &c.) = Pez. tuberosæ.  
*d.* Patellariæ (Sarea, Patellaria, Cenangium, &c.)  
*e.* Phacidiaei (Glonium, Actidium, Hysterium, Phacidium, Rhytisma, Leptostroma, &c.)  
*f.* Stictidei.
3. PYRENOMYCETES. { Genera e prorsus novo principio ultra 50, naturalissima.  
{ *a.* Sphæciei. Ostiolo prominulo, sive eustomi, quale etiam Sp. fuscæ, &c.  
{ *b.* Dothiorei. Ostiolo impresso, sive cryptostomi.  
{ *c.* Dichænei. Ostiolo rimoso, sive rhegmostomi.  
{ *d.* Perisporiei. Ostiolo poroso, sive porostomi. Analogicum Gasteromycetibus.  
{ *e.* Apiosporiei. Astomi nec sponte dehiscentes (h. l. Antennaria, Myxothecium, Vermicularia.)  
{ *f.* Cytisporiei. Coniomycetes in nucleum redacti, tangunt stilbosporeos.

ginal design. It will contain a very full account of all that has been hitherto written on the subject; and an original figure of every species and remarkable variety will be given. Dr. Johnston intends at the same time to avail himself of this opportunity of adding many new figures of zoophytes contributed by his friends, and acquired by himself, since the volume on British Zoophytes was published.

*Corpus Florarum Provincialium Sueciæ.*—I. *Floram Scanicam scripsit* E. Fries, Upsaliæ, 1835. 8vo.

This is the first of a series of Floras of the different parts of Sweden, comprising that of Schonen. Our object in noticing it here is to give a sketch of the learned Professor's latest views on the arrangement of Fungi, which we are enabled to do not only by the help of the work itself, but from a letter from the Professor himself, dated October 4th, 1838. As the work (though, like the 'Novitiæ Floræ Suecicæ' lately noticed, it is of great utility) is not probably in many hands, we think it cannot fail to be acceptable.

- |  |   |  |
|--|---|--|
| 1. HYMENOMYCETES.                                  | { | a. Agaricinei. b. Polyporei. c. Hydnei. d. Auricularinei. e. Clavariiei. f. Tremellinei.     |
| 2. DISCOMYCETES.                                   | { | a. Helvellei (Morchella, Helvella, Rhizina, Peziza, &c.) = Pez. ceracæ.                      |
| <i>Discus excipulo saturatior.</i>                 | { | b. Bulgariiei (Leotia, Bulgaria, &c.) = Pez. gelatinosæ.                                     |
|  | { | c. Dermei (Vibrissea, Ditiola, Dermea, &c.) = Pez. tuberosæ.                                 |
| <i>Discus ut in sequentibus excipulo dilutior.</i> | { | d. Patellariiei (Sarea, Patellaria, Cenangium, &c.)  |
|  | { | e. Phacidiaei (Glonium, Actidium, Hysterium, Phacidium, Rhytisma, Leptostroma, &c.)          |
|  | { | f. Stictidei.  |
| 3. PYRENOMYCETES.                                  | { | Genera e prorsus novo principio ultra 50, naturalissima.                                     |
|  | { | a. Sphæciei. Ostiolo prominulo, sive eustomi, quale etiam Sp. fuscæ, &c.                     |
|  | { | b. Dothiorei. Ostiolo impresso, sive cryptostomi.  |
|  | { | c. Dichænei. Ostiolo rimoso, sive rhegmostomi.   |
|  | { | d. Perisporiei. Ostiolo poroso, sive porostomi. Analogicum Gasteromycetibus.                 |
|  | { | e. Apiosporiei. Astomi nec sponte dehiscentes (h. l. Antennaria, Myxothecium, Vermicularia.) |
|  | { | f. Cytisporiei. Coniomycetes in nucleum redacti, tangunt stilbosporeos.                      |

ginal design. It will contain a very full account of all that has been hitherto written on the subject; and an original figure of every species and remarkable variety will be given. Dr. Johnston intends at the same time to avail himself of this opportunity of adding many new figures of zoophytes contributed by his friends, and acquired by himself, since the volume on British Zoophytes was published.

*Corpus Florarum Provincialium Sueciæ.*—I. *Floram Scanicam scripsit* E. Fries, Upsaliæ, 1835. 8vo.

This is the first of a series of Floras of the different parts of Sweden, comprising that of Schonen. Our object in noticing it here is to give a sketch of the learned Professor's latest views on the arrangement of Fungi, which we are enabled to do not only by the help of the work itself, but from a letter from the Professor himself, dated October 4th, 1838. As the work (though, like the 'Novitiæ Floræ Suecicæ' lately noticed, it is of great utility) is not probably in many hands, we think it cannot fail to be acceptable.

1. HYMENOMYCETES. { *a.* Agaricinei. *b.* Polyporei. *c.* Hydnei. *d.* Auricularinei. *e.* Clavariæ. *f.* Tremellinei.
2. DISCOMYCETES. { *a.* Helvellei (Morchella, Helvella, Rhizina, Peziza, &c.) = Pez. ceracæ.  
*Discus excipulo saturatior.* { *b.* Bulgariæ (Leotia, Bulgaria, &c.) = Pez. gelatinosæ.  
*Discus ut in sequentibus excipulo dilutior.* { *c.* Dermei (Vibrissea, Ditiola, Dermea, &c.) = Pez. tuberosæ.  
*d.* Patellariæ (Sarea, Patellaria, Cenangium, &c.)  
*e.* Phacidiaei (Glonium, Actidium, Hysterium, Phacidium, Rhytisma, Leptostroma, &c.)  
*f.* Stictidei.
3. PYRENOMYCETES. { Genera e prorsus novo principio ultra 50, naturalissima.  
{ *a.* Sphæciei. Ostiolo prominulo, sive eustomi, quale etiam Sp. fuscæ, &c.  
{ *b.* Dothiorei. Ostiolo impresso, sive cryptostomi.  
{ *c.* Dichænei. Ostiolo rimoso, sive rhegmostomi.  
{ *d.* Perisporiei. Ostiolo poroso, sive porostomi. Analogicum Gasteromycetibus.  
{ *e.* Apiosporiei. Astomi nec sponte dehiscentes (h. l. Antennaria, Myxothecium, Vermicularia.)  
{ *f.* Cytisporiei. Coniomycetes in nucleum redacti, tangunt stilbosporeos.

ginal design. It will contain a very full account of all that has been hitherto written on the subject; and an original figure of every species and remarkable variety will be given. Dr. Johnston intends at the same time to avail himself of this opportunity of adding many new figures of zoophytes contributed by his friends, and acquired by himself, since the volume on British Zoophytes was published.

*Corpus Florarum Provincialium Sueciæ.*—I. *Floram Scanicam scripsit* E. Fries, Upsaliæ, 1835. 8vo.

This is the first of a series of Floras of the different parts of Sweden, comprising that of Schonen. Our object in noticing it here is to give a sketch of the learned Professor's latest views on the arrangement of Fungi, which we are enabled to do not only by the help of the work itself, but from a letter from the Professor himself, dated October 4th, 1838. As the work (though, like the 'Novitiæ Floræ Suecicæ' lately noticed, it is of great utility) is not probably in many hands, we think it cannot fail to be acceptable.

1. HYMENOMYCETES. { *a.* Agaricinei. *b.* Polyporei. *c.* Hydnei. *d.* Auricularinei. *e.* Clavariæ. *f.* Tremellinei.
2. DISCOMYCETES. { *a.* Helvellei (Morchella, Helvella, Rhizina, Peziza, &c.) = Pez. ceracæ.  
*Discus excipulo saturatior.* { *b.* Bulgariæ (Leotia, Bulgaria, &c.) = Pez. gelatinosæ.  
*Discus ut in sequentibus excipulo dilutior.* { *c.* Dermei (Vibrissea, Ditiola, Dermea, &c.) = Pez. tuberosæ.  
*d.* Patellariæ (Sarea, Patellaria, Cenangium, &c.)  
*e.* Phacidiaei (Glonium, Actidium, Hysterium, Phacidium, Rhytisma, Leptostroma, &c.)  
*f.* Stictidei.
3. PYRENOMYCETES. { Genera e prorsus novo principio ultra 50, naturalissima.  
{ *a.* Sphæciei. Ostiolo prominulo, sive eustomi, quale etiam Sp. fuscæ, &c.  
{ *b.* Dothiorei. Ostiolo impresso, sive cryptostomi.  
{ *c.* Dichænei. Ostiolo rimoso, sive rhegmostomi.  
{ *d.* Perisporiei. Ostiolo poroso, sive porostomi. Analogicum Gasteromycetibus.  
{ *e.* Apiosporiei. Astomi nec sponte dehiscentes (h. l. Antennaria, Myxothecium, Vermicularia.)  
{ *f.* Cytisporiei. Coniomycetes in nucleum redacti, tangunt stilbosporeos.

4. GASTEROMYCETES.  $\left\{ \begin{array}{l} a. \text{ Phalloidei. Analogi cum Hymenomycetibus.} \\ b. \text{ Tuberacei. Analogi cum Discomycetibus.} \\ c. \text{ Nidularacei. Analogi cum Pyrenomycetibus.} \\ d. \text{ Lycoperdinei.} \\ e. \text{ Trichodermacei. Hyphomycetes perfectius evolu-} \\ \text{ luti.} \\ f. \text{ Myxogastres. Hypodermiis eximie analogi v. c.} \\ \text{ Perichæna strobilina = Peridermium, Lk.} \end{array} \right.$
- Sporidia fixa.*
- Sporidia pulveracea laxa.*
5. HYPHOMYCETES.  $\left\{ \begin{array}{l} (\text{Obs. nunc limites non e substantia floccosa, sed in} \\ \text{ morphosi pono.}) \\ a. \text{ Isariei sive Hymenomycetes elementares.} \\ b. \text{ Stilbini sive Discomycetes elementares.} \\ c. \text{ Mucorini sive Pyrenomycetes.} \\ d. \text{ Mucedines sive Gasteromycetes elementares cum} \\ \text{ Trichodermaceis confluentes.} \\ e. \text{ Dematiei.} \\ f. \text{ Sporodermei sive Coniomycetes in floccos conca-} \\ \text{ tenati.} \end{array} \right.$
6. CONIOMYCETES.  $\left\{ \begin{array}{l} a. \text{ Sclerotiacei sive Hymenomycetes elementares.} \\ b. \text{ Tubercularei sive Discomycetes elementares.} \\ c. \text{ Stilbosporei sive Pyrenomycetes elementares.} \\ d. \text{ Illosporei sive Gasteromycetes elementares.} \\ e. \text{ Sepedonie sive Hyphomycetes elementares.} \\ f. \text{ Hypodermiei.} \end{array} \right.$

## PROCEEDINGS OF LEARNED SOCIETIES.

## ZOOLOGICAL SOCIETY.

February 12, 1839.—Thomas Bell, Esq., V.P., in the Chair.

A paper from Mr. Schomburgk, entitled "Remarks on the Greater Ant-bear (*Myrmecophaga jubata*)," was read. This paper commences with some general observations on the *Edentata* and *Monotremata*: the author then proceeds to give a detailed description of the animal under consideration. The following is an abstract of the remaining portion of the paper, or that which relates to the habits of the animal.

Mr. Schomburgk observes, that at a distance the Ant-bear appears to be a much taller animal than it really is, owing to the elongated and nearly erect hair of the mane, and also the erect manner in which it carries its large bushy tail. When walking, the outer portion of the fore foot is applied to the ground, and the long claws are then doubled inwards. It runs with a peculiar trot, and is not, as has been represented, slow in its movements and easily overtaken; for when chased it will keep a horse in canter, and does not tire readily.

4. GASTEROMYCETES.  $\left\{ \begin{array}{l} a. \text{ Phalloidei. Analogi cum Hymenomycetibus.} \\ b. \text{ Tuberacei. Analogi cum Discomycetibus.} \\ c. \text{ Nidularacei. Analogi cum Pyrenomycetibus.} \\ d. \text{ Lycoperdinei.} \\ e. \text{ Trichodermacei. Hyphomycetes perfectius evolu-} \\ \text{ luti.} \\ f. \text{ Myxogastres. Hypodermiis eximie analogi v. c.} \\ \text{ Perichæna strobilina = Peridermium, Lk.} \end{array} \right.$
- Sporidia fixa.*
- Sporidia pulveracea laxa.*
5. HYPHOMYCETES.  $\left\{ \begin{array}{l} (\text{Obs. nunc limites non e substantia floccosa, sed in} \\ \text{ morphosi pono.}) \\ a. \text{ Isariei sive Hymenomycetes elementares.} \\ b. \text{ Stilbini sive Discomycetes elementares.} \\ c. \text{ Mucorini sive Pyrenomycetes.} \\ d. \text{ Mucedines sive Gasteromycetes elementares cum} \\ \text{ Trichodermaceis confluentes.} \\ e. \text{ Dematiei.} \\ f. \text{ Sporodermei sive Coniomycetes in floccos conca-} \\ \text{ tenati.} \end{array} \right.$
6. CONIOMYCETES.  $\left\{ \begin{array}{l} a. \text{ Sclerotiacei sive Hymenomycetes elementares.} \\ b. \text{ Tubercularei sive Discomycetes elementares.} \\ c. \text{ Stilbosporei sive Pyrenomycetes elementares.} \\ d. \text{ Illosporei sive Gasteromycetes elementares.} \\ e. \text{ Sepedoniei sive Hyphomycetes elementares.} \\ f. \text{ Hypodermiei.} \end{array} \right.$

## PROCEEDINGS OF LEARNED SOCIETIES.

## ZOOLOGICAL SOCIETY.

February 12, 1839.—Thomas Bell, Esq., V.P., in the Chair.

A paper from Mr. Schomburgk, entitled “Remarks on the Greater Ant-bear (*Myrmecophaga jubata*),” was read. This paper commences with some general observations on the *Edentata* and *Monotremata*: the author then proceeds to give a detailed description of the animal under consideration. The following is an abstract of the remaining portion of the paper, or that which relates to the habits of the animal.

Mr. Schomburgk observes, that at a distance the Ant-bear appears to be a much taller animal than it really is, owing to the elongated and nearly erect hair of the mane, and also the erect manner in which it carries its large bushy tail. When walking, the outer portion of the fore foot is applied to the ground, and the long claws are then doubled inwards. It runs with a peculiar trot, and is not, as has been represented, slow in its movements and easily overtaken; for when chased it will keep a horse in canter, and does not tire readily.

4. GASTEROMYCETES.  $\left\{ \begin{array}{l} a. \text{ Phalloidei. Analogi cum Hymenomycetibus.} \\ b. \text{ Tuberacei. Analogi cum Discomycetibus.} \\ c. \text{ Nidularacei. Analogi cum Pyrenomycetibus.} \\ d. \text{ Lycoperdinei.} \\ e. \text{ Trichodermacei. Hyphomycetes perfectius evolu-} \\ \text{ luti.} \\ f. \text{ Myxogastres. Hypodermiis eximie analogi v. c.} \\ \text{ Perichæna strobilina = Peridermium, Lk.} \end{array} \right.$
- Sporidia fixa.*
- Sporidia pulveracea laxa.*
5. HYPHOMYCETES.  $\left\{ \begin{array}{l} (\text{Obs. nunc limites non e substantia floccosa, sed in} \\ \text{ morphosi pono.}) \\ a. \text{ Isariei sive Hymenomycetes elementares.} \\ b. \text{ Stilbini sive Discomycetes elementares.} \\ c. \text{ Mucorini sive Pyrenomycetes.} \\ d. \text{ Mucedines sive Gasteromycetes elementares cum} \\ \text{ Trichodermaceis confluentes.} \\ e. \text{ Dematiei.} \\ f. \text{ Sporodermei sive Coniomycetes in floccos conca-} \\ \text{ tenati.} \end{array} \right.$
6. CONIOMYCETES.  $\left\{ \begin{array}{l} a. \text{ Sclerotiacei sive Hymenomycetes elementares.} \\ b. \text{ Tubercularei sive Discomycetes elementares.} \\ c. \text{ Stilbosporei sive Pyrenomycetes elementares.} \\ d. \text{ Illosporei sive Gasteromycetes elementares.} \\ e. \text{ Sepedoniei sive Hyphomycetes elementares.} \\ f. \text{ Hypodermiei.} \end{array} \right.$

## PROCEEDINGS OF LEARNED SOCIETIES.

## ZOOLOGICAL SOCIETY.

February 12, 1839.—Thomas Bell, Esq., V.P., in the Chair.

A paper from Mr. Schomburgk, entitled “Remarks on the Greater Ant-bear (*Myrmecophaga jubata*),” was read. This paper commences with some general observations on the *Edentata* and *Monotremata*: the author then proceeds to give a detailed description of the animal under consideration. The following is an abstract of the remaining portion of the paper, or that which relates to the habits of the animal.

Mr. Schomburgk observes, that at a distance the Ant-bear appears to be a much taller animal than it really is, owing to the elongated and nearly erect hair of the mane, and also the erect manner in which it carries its large bushy tail. When walking, the outer portion of the fore foot is applied to the ground, and the long claws are then doubled inwards. It runs with a peculiar trot, and is not, as has been represented, slow in its movements and easily overtaken; for when chased it will keep a horse in canter, and does not tire readily.

4. GASTEROMYCETES.  $\left\{ \begin{array}{l} a. \text{ Phalloidei. Analogi cum Hymenomycetibus.} \\ b. \text{ Tuberacei. Analogi cum Discomycetibus.} \\ c. \text{ Nidularacei. Analogi cum Pyrenomycetibus.} \\ d. \text{ Lycoperdinei.} \\ e. \text{ Trichodermacei. Hyphomycetes perfectius evolu-} \\ \text{ luti.} \\ f. \text{ Myxogastres. Hypodermiis eximie analogi v. c.} \\ \text{ Perichæna strobilina = Peridermium, Lk.} \end{array} \right.$
- Sporidia fixa.*
- Sporidia pulveracea laxa.*
5. HYPHOMYCETES.  $\left\{ \begin{array}{l} (\text{Obs. nunc limites non e substantia floccosa, sed in} \\ \text{ morphosi pono.}) \\ a. \text{ Isariei sive Hymenomycetes elementares.} \\ b. \text{ Stilbini sive Discomycetes elementares.} \\ c. \text{ Mucorini sive Pyrenomycetes.} \\ d. \text{ Mucedines sive Gasteromycetes elementares cum} \\ \text{ Trichodermaceis confluentes.} \\ e. \text{ Dematiei.} \\ f. \text{ Sporodermei sive Coniomycetes in floccos conca-} \\ \text{ tenati.} \end{array} \right.$
6. CONIOMYCETES.  $\left\{ \begin{array}{l} a. \text{ Sclerotiacei sive Hymenomycetes elementares.} \\ b. \text{ Tubercularei sive Discomycetes elementares.} \\ c. \text{ Stilbosporei sive Pyrenomycetes elementares.} \\ d. \text{ Illosporei sive Gasteromycetes elementares.} \\ e. \text{ Sepedoniei sive Hyphomycetes elementares.} \\ f. \text{ Hypodermiei.} \end{array} \right.$

## PROCEEDINGS OF LEARNED SOCIETIES.

## ZOOLOGICAL SOCIETY.

February 12, 1839.—Thomas Bell, Esq., V.P., in the Chair.

A paper from Mr. Schomburgk, entitled "Remarks on the Greater Ant-bear (*Myrmecophaga jubata*)," was read. This paper commences with some general observations on the *Edentata* and *Monotremata*: the author then proceeds to give a detailed description of the animal under consideration. The following is an abstract of the remaining portion of the paper, or that which relates to the habits of the animal.

Mr. Schomburgk observes, that at a distance the Ant-bear appears to be a much taller animal than it really is, owing to the elongated and nearly erect hair of the mane, and also the erect manner in which it carries its large bushy tail. When walking, the outer portion of the fore foot is applied to the ground, and the long claws are then doubled inwards. It runs with a peculiar trot, and is not, as has been represented, slow in its movements and easily overtaken; for when chased it will keep a horse in canter, and does not tire readily.